arXiv:1108.1265v1 [math.GT] 5 Aug 2011

WRINKLED EMBEDDINGS

To Paul Schweitzer on his 70th birthday

Y. M. Eliashberg *
Stanford University,
Stanford, CA 94305 USA

N. M. Mishachev
Lipetsk Technical University,
Lipetsk, 398055 Russia

Abstract

A *wrinkled embedding* $f : V^n \to W^m$ is a topological embedding which is a smooth embedding everywhere on V except a set of $(n-1)$-dimensional spheres, where f has cuspidal corners. In this paper we prove that any rotation of the tangent plane field $TV \subset TW$ of a *smoothly embedded* submanifold $V \subset W$ can be approximated by a homotopy of *wrinkled embeddings* $V \to W$.

Contents

1 Introduction **2**
1.1 Wrinkled embeddings 2
1.2 Main result and the idea of the proof 3
1.3 Applications 4
1.4 History of the problem 4
1.5 Remark 5

2 Integrable approximations of tangential homotopies **5**
2.1 Tangential homotopies of embeddings 5
2.2 Local integrable approximations 6
2.3 Wrinkled embeddings 7
2.4 Fibered wrinkled embeddings 10

*Partially supported by the NSF grant DMS-0707103

2.5 Main theorem 12
2.6 Local integrable approximations near wrinkles 13
2.7 Main lemma 14
2.8 Proof of Theorem 2.5.2 20
2.9 Fibered case 21
2.10 Double folds 22

3 Applications **24**
3.1 Homotopy principle for directed wrinkled embeddings 24
3.2 Embeddings into foliations 24
3.3 Embeddings into distributions 29

4 Appendix: Wrinkling **30**
4.1 Folds and cusps 31
4.2 Wrinkles and wrinkled mappings 32
4.3 Fibered wrinkles and fibered wrinkled mappings 33
4.4 Main theorems about wrinkled mappings 33
4.5 Wrinkled mappings into foliations 34
4.6 Generalized wrinkled mappings 36

1 Introduction

1.1 Wrinkled embeddings

A *wrinkled embedding* $f : V^n \to W^m$, $n < m$, is a topological embedding which is a smooth embedding everywhere on V except a finite set of $(n-1)$-dimensional spheres $S_i \subset V$, where f has cuspidal corners: threefold corners along an equator $S_i' \subset S_i$ and twofold corners along the complement $S_i \setminus S_i'$. For $n = 2$ and $q = 3$ see Fig. 1 and Fig. 2. The spheres S_i are called *wrinkles*. A formal definition of wrinkled embeddings is given in Section 2.3 below.
Note that for $n = 1$ each wrinkle consists of two twofold cuspidal points. *Families* of wrinkled embeddings may have, in addition to wrinkles, *embryos* of wrinkles, and therefore wrinkles may *appear* and *disappear* in a homotopy of wrinkled embeddings.
Let us point out that a *wrinkled embedding is not a wrinkled map* in the sense of [EM97]. The relation between the two notions is discussed in Section 2.3 below.



Figure 1: The wrinkled embedding $f : V^2 \to W^3$ near $S_i \setminus S_i'$



Figure 2: The wrinkled embedding $f : V^2 \to W^3$ near S_i'

1.2 Main result and the idea of the proof

In this paper we prove (see Theorem 2.5.1 below) that any homotopy of the tangent plane field $TV \subset TW$ (*tangential homotopy*) of a *smoothly embedded* submanifold $V \subset W$ can be approximated by a *wrinkled isotopy*, i.e. an isotopy through wrinkled embeddings $V \to W$. For $n = 1$ the idea of the proof is presented on Fig. 3. Here we consider the counterclockwise tangential rotation of an interval in $\mathbb{R}^2$.



Figure 3: Wrinkled approximation of a counterclockwise tangential rotation

The implementation of this general idea for $n > 1$ is far from being straightforward. There are lot of similarities here with the Nash-Kuiper theorem about isometric C^1-embeddings $V^n \to W^{n+1}$, where the proof in the case $n = 1$ is more or less trivial, while already contains the general idea (goffering). However, its realization for $n > 1$ is highly non-trivial.

1.3 Applications

Our main theorem can be reformulated as an *h-principle for A-directed wrinkled embeddings*, see 3.1.1. As an application of the main theorem we prove an h-principle for *embeddings* $f : V^n \to (W, \xi)$, $n \geqslant q = \operatorname{codim} \xi$, whose tangency singularities with respect to a distribution ξ (integrable or non-integrable) are simple, e.g. folds, or alternatively *generalized* wrinkles (see Theorem 3.2.1 and Section 3.3). This h-principle allows us, in particular, to simplify the singularities of an individual embedding or a family of embeddings $V^n \to \mathbb{R}^m$, $n < m$, with respect to the projection $\mathbb{R}^m = \mathbb{R}^q \times \mathbb{R}^{m-q} \to \mathbb{R}^q$, $n \geqslant q$, i.e with respect to the standard foliation of $\mathbb{R}^m$ by $(m-q)$-dimensional affine subspaces, parallel to $0 \times \mathbb{R}^{m-q}$.

1.4 History of the problem

A. Directed embeddings. An embedding $V^n \to W^m$ is called *A-directed*, if its tangential (or Gaussian) image belongs to a given subset A of the Grassmannian bundle $\mathrm{Gr}_n W$. Using his convex integration method, Gromov proved in [Gr86] a general h-principle for A-directed embedding in the case when V is an *open* manifold and $A \subset \mathrm{Gr}_n W$ is an open subset. C. Rourke and B. Sanderson gave two independent proofs of this theorem in [RS01]. A different proof based on our holonomic approximation theorem was given in our book [EM02]. For some special $A \subset \mathrm{Gr}_n W$ Gromov also proved in [Gr73] and [Gr86] the h-principle for *closed* manifolds. However, for closed V the h-principle for A-directed embeddings fails for a *general* open A. For example, for any closed V^n there is no A-directed embeddings $V^n \to \mathbb{R}^{n+1}$ unless $\pi(A) = S^n$, where

$$\pi : \mathrm{Gr}_n \mathbb{R}^{n+1} = S^n \times \mathbb{R}^{n+1} \to S^n$$

is the projection. The main theorem of the current paper states that *this h-principle can be saved by relaxing the notion of embedding.* D. Spring in

[Sp05] proved, using Gromov's convex integration method and the *geometry of spiral curves* from [Sp02], an existence theorem for directed embeddings with *twofold spherical corners*, which is equivalent to the non-parametric version of our main theorem, see the discussion in Sections 2.10 and 3.1 below.

B. Simplification of singularities. First result allowing to simplify singularities of an *individual* embedding $V^n \to \mathbb{R}^m$, $n < m$, with respect to the projection $\mathbb{R}^m = \mathbb{R}^q \times \mathbb{R}^{m-q} \to \mathbb{R}^q$, $n \geqslant q$, was proven in [El72]. In fact, in [El72] this result was formulated for $n = m - 1$; the general case $n < m$ can be derived from this basic one by induction which was done in [EM00]. A different proof based on convex integration method was given by D. Spring in [Sp02]. In [EM00] we also proved the *parametric* version, but only for $q = 1$, and only the epimorphism part of the corresponding parametric h-principle. However, the approach in [EM00] does not seem to be suitable to recover the main results of the current paper.
The reader may find additional interesting information related to the subject of this paper in [Sp02] and [Sp05]. A different approach to the problem of simplification of singularities can be found in [RS03].

1.5 Remark

We assume that the reader is familiar with the general philosophy of the h-principle, see [Gr86] and [EM02]. It is useful, though not necessary for the reader to be also familiar with the Holonomic Approximation Theorem from [EM02]) and the wrinkling philosophy (see [EM97] and [EM98]). We recall in Section 4, for a convenience of the reader, some definitions and results from [EM97] and [EM98] and introduce there the notions of *generalized wrinkles* and *generalized wrinkled maps*.

2 Integrable approximations of tangential homotopies

2.1 Tangential homotopies of embeddings

In what follows we assume that $V \subset W$ is an embedded compact submanifold and denote by f_0 the inclusion $\mathrm{i}_V : V \hookrightarrow W$. We also assume that the manifolds W and $\mathrm{Gr}_n W$ are endowed with Riemannian metrics.

Let $\pi : \mathrm{Gr}_n W \to W$ be the Grassmannian bundle of tangent n-planes to a m-dimensional manifold W, $m > n$, and V a n-dimensional manifold. Given a *monomorphism* (= fiberwise injective homomorphism) $F : TV \to TW$, we will denote by $\mathrm{G}F$ the corresponding map $V \to \mathrm{Gr}_n W$. Thus the tangential (Gaussian) map associated with an immersion $f : V \to W$ can be written as $\mathrm{G}df$.

A *tangential homotopy* of an embedding f_0 is a homotopy $G_t : V \to \mathrm{Gr}_n W$, such that $G_0 = \mathrm{G}df_0$ which covers an isotopy $g_t = \pi \circ G_t : V \to W$, $g_0 = f_0$. A tangential homotopy $G_t : V \to \mathrm{Gr}_n W$ is called *integrable*, if $G_t = \mathrm{G}dg_t$. A tangential homotopy $G_t : V \to \mathrm{Gr}_n W$ is called a *tangential rotation*, if $G_0 = \mathrm{G}df_0$ and $\pi \circ G_t = f_0$, i.e. G_t covers the constant isotopy $g_t = f_0 : V \to W$.

◀ **Problem.** Let $G_t : V \to \mathrm{Gr}_n W$ be a tangential homotopy. We want to construct an arbitrarily close *integrable approximation* of G_t, i.e an isotopy of embeddings $f_t : V \to W$, such that $\mathrm{G}df_t$ is arbitrarily close to G_t. One can reduce the problem to the case when G_t is a *tangential rotation*. Indeed, we can consider, instead of G_t, the rotation $(d\hat{g}_t)^{-1} \circ G_t$, where $\hat{g}_t : W \to W$ is a diffeotopy which extends the isotopy $g_t = \pi \circ G_t : V \to W$. Note that if G_t is an *integrable* tangential homotopy, then $(d\hat{g}_t)^{-1} \circ G_t$ is the constant homotopy $G_t = G_0$. ▶

Of course, an integrable approximation of a tangential rotation does not exist in general. However, as we shell see, one can always construct an integrable approximation of G_t by a family of *wrinkled embeddings*.

2.2 Local integrable approximations

Let $X \subset W$ be a tubular neighborhood of $V \subset W$, and $\pi : X \to V$ the normal projection. Denote by $\mathcal{N}$ the normal foliation of X by the fibers of π. An isotopy $f_t : V \to W$, $f_0 = i_V$, is called *graphical*, if all the images $f_t(V)$ are transversal to $\mathcal{N}$. In other words, the graphical isotopy $V \to W$ is an isotopy of sections $V \to X$, up to reparameterizatons of V. A tangential rotation $G_t : V \to \mathrm{Gr}_n W$ is called *small*, if $G_t(v)$ is transversal to $\mathcal{N}$ for all t and v.

Following Gromov's book [Gr86] we will use the notation $\mathcal{O}p\, A$ as a replacement of the expression *an open neighborhood of* $A \subset V$. In other words, $\mathcal{O}p\, A$ is an arbitrarily small but not specified open neighborhood of a subset $A \subset V$.

2.2.1. (Local integrable approximation of a small tangential rotation) *Let $G_t : V \to \mathrm{Gr}_n W$ be a small tangential rotation of $V \subset W$ and $K \subset V$ a stratified subset of* **positive** *codimension. Then there exists an arbitrarily C^0-small graphical isotopy $f_t : V \to W$ such that the homotopy*

$$\mathrm{G}df_t|_{\mathcal{O}p\, K} : \mathcal{O}p\, K \to \mathrm{Gr}_n W$$

is arbitrarily C^0-close to the tangential rotation $G_t|_{\mathcal{O}p\, K}$.

Proof. The space $X^{(1)}$ of 1-jets of sections $V \to X$ can be interpreted as a space of tangent to X n-planes which are non-vertical, i.e. transversal to $\mathcal{N}$. Hence the inclusion $f_0 : V \hookrightarrow X$ together with the tangential homotopy $G_t : V \to \mathrm{Gr}_n W$ can be viewed as a homotopy of sections $F_t : V \to X^{(1)}$. For arbitrarily small ε and δ we can construct, using Holonomic Approximation Theorem 3.1.2 in [EM02], a holonomic ε-approximation $\widetilde{F}_t$ of F_t over $\mathcal{O}p\, \widetilde{h}_t(K)$, where $\widetilde{h}_t : V \to V$ is a δ-small diffeotopy. The 0-jet part $\widetilde{f}_t$ of the section $\widetilde{F}_t$ is automatically an embedding, because $\widetilde{f}_t$ is a section of the normal bundle. Thus, we have a family $\widetilde{f}_t$ of integrable approximations of G_t over $\mathcal{O}p\, \widetilde{h}_t(K)$, and hence one can define the required isotopy $f_t : V \to W$ as the composition $\widetilde{f}_t \circ \widetilde{h}_t$, where $\widetilde{f}_t : V \to X$ is an extension of the isotopy $\widetilde{f}_t : \mathcal{O}p\, \widetilde{h}^t(K) \to X$ to an isotopy of *sections* $V \to X$. □

◀ **Remarks. 1.** One can also apply the above construction to *any* section $\widetilde{V} \subset X$ instead of the zero-section $V \subset X$, provided that the tangential rotation of $T\widetilde{V}$ is transversal to $\mathcal{N}$ (here $\mathcal{N}$ is the *original* foliation on X by the fibers of the projection $\pi : X \to V$).

2. Without the "graphical" restriction on f_t Theorem 2.2.1 remains true for *any* tangential rotation, see 4.4.1 in [EM02].

3. The relative and the parametric versions of the holonomic approximation theorem similarly prove the relative and the parametric versions of Theorem 2.2.1. In the relative version the homotopy G_t and the diffeotopy h_t are constant near a compact subset $L \subset K$. In the parametric version we deal with a family of embedded submanifolds $V_p \hookrightarrow W$, $p \in B$. The parameters space B usually is a manifold, possibly with non-empty boundary ∂B, and the homotopies $G_{t,p}$, $p \in B$, are constant for $p \in \mathcal{O}p\, \partial B$. In this case the diffeotopies $h_{t,p}$, $p \in B$, are constant for $p \in \mathcal{O}p\, \partial B$. ▶

2.3 Wrinkled embeddings

A. Definition. A (smooth) map $f : V^n \to W^m$, $n < m$, is called a *wrinkled*

embedding, if

- f is a topological embedding;
- any connected component S_i of the singularity $S = \Sigma(f)$ is diffeomorphic to the standard $(n-1)$-dimensional sphere S^{n-1} and bounds an n-dimensional disk $D_i \subset V$;
- the map f near each sphere S_i is equivalent to the map

$$Z(n,m) : \mathcal{O}p_{\,\mathbb{R}^n} S^{n-1} \to \mathbb{R}^m$$

given by the formula

$$(y_1, ..., y_{n-1}, z) \longmapsto$$

$$\left(y_1, ..., y_{n-1}, z^3 + 3(|y|^2 - 1)z, \int_0^z (z^2 + |y|^2 - 1)^2 dz,\, 0, ..., 0\right). \qquad (1)$$

The compositions of the canonical form (1) with the projection to the space of first n local coordinates in $\mathbb{R}^m$ is the standard equidimensional wrinkled map $\mathbb{R}^n \to \mathbb{R}^n$, see Section 4.2 or [EM97]. Thus, the canonical form for the wrinkled embeddings contains, in comparison to the canonical form for the wrinkled mappings, the *unfolding* function $\int_0^z (z^2 + |y|^2 - 1)^2\, dz$ and zero functions as additional coordinates in the image. The spheres S_i and its images $f(S_i)$ are called *wrinkles* of the wrinkled embedding f. According to the formula (1), each wrinkle S_i has a marked $(n-2)$-dimensional equator $S_i' \subset S_i$ such that

- the local model for f near each point of $S_i \setminus S_i'$ is given by the formula

$$(y_1, ..., y_{n-1}, z) \longmapsto (y_1, ..., y_{n-1}, z^2, z^3, 0, ..., 0)\,, \qquad (2)$$

see Fig. 1;

- the local model for f near each point of S_i' is given by the formula

$$(y_1, ..., y_{n-1}, z) \longmapsto (y_1, ..., y_{n-1}, z^3 - 3y_1 z, \int_0^z (z^2 - y_1)^2 dz, 0, ..., 0), \quad (3)$$

see Fig. 2.

The compositions of the canonical forms (2) and (3) with the projection to the space of first n local coordinates in $\mathbb{R}^m$ are the standard fold and cusp of the equidimensional map $\mathbb{R}^n \to \mathbb{R}^n$.
We will denote the union $\bigcup\limits_i S'_i$ of all the equators S'_i by S'.

The restriction of a wrinkled embedding f to $\operatorname{Int} D_i$ *may have singularities*, which are, of course, again wrinkles. We will say, that a wrinkle S_i *has a depth* $d+1$, if S_i is contained in d "exterior" wrinkles. The depth of a wrinkled embedding is the maximal depth of its wrinkles; according to this definition the smooth embeddings are the wrinkled embeddings of depth 0.
◀ **Remark.** Note that a wrinkled embedding f near $S \setminus S'$ is equivalent to the restriction of a generic Σ^{11}-map of an $(n+1)$-dimensional manifold to its fold Σ^1 near the cusp Σ^{11}. However, near S' a wrinkled embedding f *is not equivalent* to the restriction of a generic Σ^{111}-map to Σ^1 near the swallow tail singularity Σ^{111}. ▶

B. Regularization. Any wrinkled embedding can be canonically *regularized* by changing the unfolding function $u(y,z) = \int (z^2 + |y|^2 - 1)^2\, dz$ in the canonical form to a C^1-close function $\widetilde{u}(y,z)$ such that $\partial_z \widetilde{u}(y,z) > 0$, see Fig. 4 and Fig. 5. One can chose $\widetilde{u}$ such that $\widetilde{u}(y,z) = u(y,z)$ for all $(y,z) \in S \setminus \mathcal{O}p\, S'$. Then the respective regularization does not moves the twofold corner points $f(S \setminus S')$ everywhere except an arbitrarily small neighborhood of the threefold corner points $f(S')$.



Figure 4: Regularization of a wrinkled embedding near $\Sigma_i \setminus \Sigma'_i$

C. Gaussian map.
For any wrinkled embedding $f : V \to W$ and any $v_0 \in \Sigma_f$ there exists a limit

$$\lim_{v \to v_0,\ v \in V \setminus \Sigma_f} f_*(T_v V).$$

Hence, we can associate with a wrinkled embedding its *wrinkled tangent bundle* $T(f)$. If V is oriented then the bundle $T(f)$ is oriented as well. The



Figure 5: Regularization of a wrinkled embedding near Σ'_i

orientation of a tangent space T_v to V at a smooth point $v \in V \setminus \Sigma(f)$ as a fiber of $T(f)$ coincides with, or opposite to its orientation as the fiber of TV, depending on the parity of of the number of exterior wrinkles which surround v. Therefore one can canonically define a Gaussian map $\mathrm{G}df : V \to \mathrm{Gr}_n W$ into the Grassmannian of non-oriented n-planes, and if V is oriented the map $\widetilde{\mathrm{G}}df : V \to \widetilde{\mathrm{Gr}}_n W$ into the Grassmannian $\widetilde{\mathrm{Gr}}_n W$ of *oriented* n-planes in TW.

D. Normal foliation. Though the image the $f(V) \subset W$ of a wrinkled embedding $f : V \to W$ is not a smooth submanifold, one can still define an analog of normal foliation.

There exists an n-dimensional submanifold $\widehat{V} \subset W$, such that $f(\Sigma) \subset \widehat{V}$ and $\widehat{V}$ is tangent to $f(V)$ along $f(\Sigma)$. Let $X \subset W$ be a small neighborhood of $f(V)$. We supply X with an "almost normal" (to $f(V)$) foliation $\mathcal{N}$, which coincides with the normal foliations to $\widehat{V}$ near $f(\Sigma)$ and with the normal foliation to $f(V \setminus \mathcal{O}p\,\Sigma)$ near $f(V \setminus \mathcal{O}p\,\Sigma)$.

2.4 Fibered wrinkled embeddings

The notion of a wrinkled embedding can be extended to the parametric case. Considering k-parametric *families* f_p, $p \in B$, of the wrinkled embeddings we allow, in addition to the wrinkles, their *embryos*. Near each embryo $v_i \in V$ the map f_p is equivalent to the map

$$Z_0(n, m) : \mathcal{O}p_{\,\mathbb{R}^n} 0 \to \mathbb{R}^m$$

given by the formula

$$(y_1, ..., y_{n-1}, z) \longmapsto \left(y_1, ..., y_{n-1}, z^3 + 3|y|^2 z, \int (z^2 + |y|^2)^2\, dz, 0, ..., 0 \right). \quad (4)$$

Thus, wrinkles may *appear and disappear* when we consider a *homotopy* of wrinkled embeddings $f_t : V \to W$.

In a more formal mode one can use, quite similar to the case of wrinkled mappings, the "fibered" terminology (see [EM97]). A *fibered* (over B) *map* is a commutative diagram



where f is a smooth map and $p\,, q$ are submersions. For the fibered map f we denote by $T_B V$ and $T_B W$ the subbundles $\operatorname{Ker} p \subset TV$ and $\operatorname{Ker} q \subset TW$. They are tangent to foliations of V and W formed by preimages $p^{-1}(b) \subset V$, $q^{-1}(b) \subset W$, $b \in B$. We will often denote a fibered map simply by $f : V \to W$ when B, p and q are implied from the context. Fibered homotopies, fibered differentials, fibered submersions, and so on can be naturally defined in the category of fibered maps. For example, the *fibered differential* $d_B f$ of a fibered map $f : V \to W$ is the restriction

$$d_B f = df|_{T_B V} : T_B V \to T_B W\,.$$

Notice that $d_B f$ itself has the structure of a map fibered over B.
Two fibered maps, $f : V \to W$ over B and $f : V' \to W'$ over B', are called *equivalent* if there exist open subsets $A \subset B$, $A' \subset B'$, $Y \subset W$, $Y' \subset W'$ with $f(V) \subset Y$, $p(V) \subset A$, $f'(V') \subset Y'$, $p'(V') \subset A'$ and diffeomorphisms $\varphi : U \to U'$, $\psi : Y \to Y'$, $s : A \to A'$ such that they form the following commutative diagram



The canonical form $Z(k+n, k+m)$, being considered as a fibered map over the space of first k coordinates, gives us the canonical form for the *fibered wrinkle* of the *fibered wrinkled embedding*. Thus, in a k-parametric family of the wrinkled embeddings each fibered wrinkle bounds a $(k+n)$-dimensional *fibered* disk, fiberwise equivalent to the standard $(k+n)$-disk in the space $\mathbb{R}^{k+n}$, fibered over $\mathbb{R}^k$. In addition, the canonical form $Z(k+n, k+m)$ over the half space $\mathbb{R}^k_-$ $(y_1 \leqslant 0)$ gives us the model for the fibered "half-wrinkles" near the boundary of B.

2.5 Main theorem

The following Theorem 2.5.1, and its fibered analog 2.9.1 in Section 2.9 below are the main results of the paper.

2.5.1. (Integrable approximation of a tangential rotation) *Let $G_t : V \to \mathrm{Gr}_n W$ be a tangential rotation of an embedding $i_V : V \hookrightarrow W$. Then there exists a homotopy of wrinkled embeddings $f_t : V \to W$, $f_0 = i_V$, such that the homotopy $\mathrm{G}df_t : V \to \mathrm{Gr}_n W$ is arbitrarily C^0-close to G_t. If the rotation G_t is fixed on a closed subset $C \subset V$, then the homotopy f_t can be chosen also fixed on C.*

◄ **Remark.** The parametric version of Theorem 2.5.1 is also true, see Section 2.9 below. ►

A small rotation $G_t : V \to \mathrm{Gr}_n W$ is called *simple*, if $G_t(v)$ is a rotation in a $(n+1)$-dimensional subspace $L_v \subset T_v W$ for every $v \in V$ and the angle of the rotation $G_t(v)$ is less than $\pi/4$. In particular, for $q = n+1$ any small rotation with the maximum angle $< \pi/4$ is simple. Any tangential rotation can be approximated by a finite sequence of simple rotations and hence Theorem 2.5.1 follows from

2.5.2. (Integrable approximation of a simple tangential rotation) *Let $G_t : V \to \mathrm{Gr}_n W$ be a simple tangential rotation of a* **wrinkled** *embedding $f_0 : V \to W$. Then there exists a homotopy of wrinkled embeddings $f_t : V \to W$, such that the homotopy $\mathrm{G}df_t : V \to \mathrm{Gr}_n W$ is arbitrarily C^0-close to G_t. If the rotation G_t is fixed on a closed subset $C \subset V$, then the homotopy f_t can be also chosen fixed on C.*

◄ **Remark.** The proof of Theorem 2.5.2 will give us the following additional information: the homotopy f_t increases the depth of f_0 at most by 1. In particular,

- if f_0 is a smooth embedding, then f_t consists of wrinkled embeddings of depth $\leqslant 1$;
- the depth of the final map f_1 in Theorem 2.5.1 is equal to the number of simple rotations in the decomposition of the rotation G_t. ►

2.6 Local integrable approximations near wrinkles

We will distinguish the notions of *homotopy of wrinkled embeddings* and their *isotopy*. A homotopy f_t, $t \in [0,1]$, of wrinkled embeddings is called an *isotopy*, if for all $t \in [0,1$ the wrinkled embedding f_t has no embryos, i.e. its wrinkles do not die, and no new wrinkles are born. Equivalently, f_t is an isotopy if $f_t = h_t \circ f_0$ where $h_t : W \to W$ is a diffeotopy.
A C^0-small *isotopy* $f_t : V \to W$, $f_0 = i_V$, of wrinkled embeddings is called *graphical*, if all the images $f_t(V)$ are transversal to the normal foliation $\mathcal{N}$ of V. As in the smooth case, the tangential rotation $G_t : V \to \mathrm{Gr}_n W$ of the wrinkled embedding $f_0 : V \to W$ is called *small*, if $G_t(v)$ is transversal to $\mathcal{N}$ for all t and v. We will reformulate now 2.2.1 for the situation, when $f_0 : V \to W$ is a *wrinkled* embedding and $K = \Sigma = \Sigma(f_0)$.

2.6.1. (Local integrable approximation of small tangential rotation near wrinkles) *Let $G_t : V \to \mathrm{Gr}_n W$ be a small tangential rotation of a wrinkled embedding $f_0 : V \to W$. Then there exists an arbitrarily C^0-small graphical isotopy of wrinkled embeddings $f_t : V \to W$ such that the homotopy*

$$\mathrm{G}df_t|_{\mathcal{O}p\,\Sigma} : \mathcal{O}p\,\Sigma \to \mathrm{Gr}_n W$$

is arbitrarily C^0-close to the tangential rotation $G_t|_{\mathcal{O}p\,\Sigma}$.

Proof. The image $f_0(\mathcal{O}p\,\Sigma) \subset W$ is not a submanifold and hence the proof of 2.2.1 formally does not work. Let $\widehat{V} \subset W$ be an n-dimensional submanifold, such that $f_0(\Sigma) \subset \widehat{V}$ and $\widehat{V}$ is tangent to $f_0(V)$ along $f_0(\Sigma)$. Let $\widehat{G}_t$ be an extension of the rotation $G_t|_\Sigma$ to a tangential rotation $\widehat{V} \to \mathrm{Gr}_n W$. Apply 2.2.1 to the pair $(\widehat{V}, f_0(\Sigma))$, the normal foliation $\mathcal{N}$ and the tangential rotation $\widehat{G}_t$; denote the produced isotopy $\widehat{V} \to W$ by $\widehat{f}_t$. Let $g_t : W \to W$ be an ambient diffeotopy for $\widehat{f}_t$, such that $g_t^* \mathcal{N} = \mathcal{N}$. Then $f_t = g_t \circ f_0$ is the desired isotopy of wrinkled embeddings on $\mathcal{O}p\,\Sigma.$, which can be then extended to the whole V as a graphical isotopy. □

2.7 Main lemma

Lemma 2.7.1, which we prove in this section, is the main ingredient in the proof of Theorem 2.5.2.

Let us denote by $\widetilde{\mathrm{G}} : S \to S^n$ the oriented Gaussian map of an oriented hypersurface $S \subset \mathbb{R}^{n+1}$. For the angle metric on S^n denote by $U_\varepsilon(s)$ the open metric ε-neighborhood of the north pole $s = (0, \dots, 1) \in S^n$.

An oriented hypersurface $S \subset \mathbb{R}^{n+1}$ is called:

- *ε-horizontal*, if $\widetilde{\mathrm{G}}(S) \subset U_\varepsilon(s) \subset S^n$;
- *graphical*, if $\widetilde{\mathrm{G}}(S) \subset U_{\pi/2}(s) \subset S^n$;
- *ε-graphical*, if $\widetilde{\mathrm{G}}(S) \subset U_{(\pi/2)+\varepsilon}(s) \subset S^n$;
- *quasi-graphical*, if $\widetilde{\mathrm{G}}(S) \subset U_\pi(s) \subset S^n$.

Similarly a wrinkled embedding $f : S \to \mathbb{R}^{n+1}$, is called *$\varepsilon$-horizontal*, if its Gaussian image $\widetilde{\mathrm{G}}(S) \subset S^n$ is contained in $U_\varepsilon(s)$.

In what follows we will often say *almost horizontal* and *almost graphical* instead of *ε-horizontal* and *ε-graphical*, assuming that ε is appropriately small.

2.7.1. (Approximation of an embedded hypersurface by almost horizontal wrinkled embeddings) *Let $S \subset \mathbb{R}^{n+1}$ be an oriented quasi-graphical hypersurface, such that S is almost horizontal near the boundary ∂S. Then there exists a C^0-approximation of the embedding $i_S : S \hookrightarrow \mathbb{R}^{n+1}$ by an almost horizontal wrinkled embedding $f : S \to \mathbb{R}^{n+1}$ of depth 1, such that f coincides with i_S near ∂S.*

Proof. Cutting from S a small neighborhood of the critical set of the function $h = x_{n+1} : S \to \mathbb{R}$, where S is already almost horizontal, we may consider this new S, and thus assume that the function h has no critical points on S. Let $\widetilde{S} \subset \mathrm{Int} S$ be a slightly smaller compact hypersurface such that $S \setminus \widetilde{S}$ is still almost horizontal. On a neighborhood $\mathcal{O}p\, S$ consider a non-vanishing vector field v which is transversal to S, defines its given co-orientation, and horizontal, i.e. tangent to the level sets of the function h. Let v^t denotes the flow of v. We may assume that for $z \in S$ the flow-line $v^t(z)$ is defined for all $|t| \leq 1$.

Consider two foliations on S: the 1-dimensional foliation $\mathcal{G}$, formed by the gradient trajectories of the function h, and the $(n-1)$-dimensional foliation $\mathcal{L}$, formed by level surfaces of the function h.

A subset $C \subset S$ is called *cylindrical*, if C equipped with $\mathcal{L}$ and $\mathcal{G}$ is diffeomorphic to $D^{n-1} \times D^1$ equipped with the standard horizontal and vertical foliations. We call C *special* if, in addition, C is almost horizontal near its top and bottom.

Fix a covering of $\widetilde{S}$ by special cylindrical sets $C_\alpha \subset S$, $\alpha = 1, ..., K$, and fix the respective parameterizations $\varphi_\alpha : D^{n-1} \times D^1 \to C_\alpha$, which send the positive direction on D^1 to the directions on $\mathcal{G}$ defined by the gradient vector field ∇h. Given a function $\tau : S \to \mathbb{R}$, $|\tau| \leq 1$, we will denote by I_S^τ a perturbation of the inclusion i_S defined by the formula

$$I_S^\tau(z) = v^{\tau(z)}(z), \quad z \in S.$$

To clarify the geometric meaning of the construction of the necessary approximation we will give first a slightly imprecise description without technical details. Let $x_i \in D^1$, $i = 1, ..., N$, be a finite set of points such that $x_1 = -1+d/2$, $x_N = 1-d/2$ and $x_{i+1}-x_i = d$. Set $w(x) = \sum_i \delta(x-x_i)-1/d$, where $\delta(x)$ is the δ-function, and define $W(x) := \int_0^x w(x)dx$ (see Fig.6). Take



Figure 6: Functions w and W

a cut-off function $\lambda : D^{n-1} \times D^1 \to \mathbb{R}_+$, equal to 1 on a slightly lesser than $D^{n-1} \times D^1$ subset $A \subset D^{n-1} \times D^1$ and equal to 0 on $\mathcal{O}p\, \partial(D^{n-1} \times D^1)$, and define a function $\tau : D^{n-1} \times D^1 \to \mathbb{R}$ by the formula $\tau(r, x) = \lambda(r, x)W(x)$ and a function $\tau_\alpha : C_\alpha \to \mathbb{R}^1$ by the formula $\tau_\alpha(c) = \gamma(\tau \circ \varphi_\alpha^{-1}(c))$, $\gamma \in \mathbb{R}_+$. For sufficiently small d and γ the map $I_S^{\tau_\alpha}$ (see Fig.7) is a good candidate for the wrinkled embedding. Take this candidate for a moment; then assuming that all the levels $h = \varphi_\alpha(x_i)$ for all C_α are distinct from each other, we can apply the described local goffering *simultaneously* for all C_α and thus



Figure 7: Local wrinkling, the case $n = 1$

get the desired global approximation of S. Unfortunately, our map $I_S^{\tau_\alpha}$ does not match the canonical form (1) and requires an additional perturbation near the singularity. Rather than doing this, we describe below the whole construction again in a slightly modified form with all the details.

First, choose a family of curves $A_t \subset \mathbb{R}^2$, $t \in \mathbb{R}$, which is given by parametric equations

$$x(u,t) = \frac{15}{8}\int\limits_0^u (u^2 - t)^2 du\,, \quad y(u,t) = -\frac{1}{2}(u^3 - 3tu).$$

The curve A_t is a graph of a continuous function $a_t : \mathbb{R} \to \mathbb{R}$ which is smooth for $t < 0$ and smooth on $\mathbb{R} \setminus \{-\sqrt{t}, \sqrt{t}\}$ for $t \geq 0$, see Fig. 8,9. Note that

(a) $a_1(\pm 1) = \pm 1$;

(b) the composition $a_t(x(u,t))$ is a smooth function.

Choose $\sigma \in (0, \frac{1}{8})$ and consider a family of odd 1-periodic functions $\mathcal{W}_{\sigma,\,t} : \mathbb{R} \to \mathbb{R}$ with the following properties:

$$\mathcal{W}_{\sigma,\,t}(x) = \begin{cases} a_t\left(\frac{x}{\sigma}\right) & \text{for } \ x \in \mathcal{O}p\,[-\sigma, \sigma] \\ 0 & \text{for } \ x = \frac{1}{2} \end{cases}$$

$$\frac{d\mathcal{W}_{\sigma,1}}{dx}(x) \begin{cases} \geqslant 3 & \text{for } \ x \in (-\sigma, \sigma) \\ \in [-2, 0) & \text{for } \ x \in [2\sigma, \frac{1}{2} - 2\sigma] \\ \leqslant -2 & \text{for } \ x \in [\sigma, 2\sigma] \end{cases}$$



Figure 8: Function a_t, $t = 1$

(see Fig. 10). Note that the composition $\mathcal{W}_{\sigma,t}(a_t(x(u,t))$ is a smooth function. Pick an $\varepsilon > 0$ and integer $N > 0$ such that the cylinders

$$\widetilde{C}_\alpha = \varphi_\alpha(D^{n-1}_{1-2\varepsilon} \times D^1_{1-1/N})$$

are still special and cover the surface $\widetilde{S}$. We will keep ε fixed while N will need to be increased to achieve the required approximation. Choose a non-strictly decreasing smooth function $\beta : [0,1] \to \mathbb{R}$ such that

$$\beta(r) = \begin{cases} 1 & r \in [0, 1-3\varepsilon]; \\ 1 - r - \varepsilon & r \in [1-2\varepsilon, 1]. \end{cases}$$

Given an integer $N > 0$, pick also a cut-off function $\lambda : [0,1] \to [0,1]$ which is equal to 1 on $[1-\rho]$ and to 0 near 1, where $\rho = \min(\frac{1}{2N}, \frac{\varepsilon}{2})$, and define a function

$$\tau = \tau_{N,\sigma} : D^{n-1} \times D^1 \to \mathbb{R}$$

by the formula

$$\tau(r,x) = \gamma\lambda(|x|)\lambda(|r|)\mathcal{W}_{\sigma,\beta(|r|)}\left(\frac{2N+1}{2}x\right), \quad (r,x) \in D^{n-1} \times D^1, \quad \gamma \in \mathbb{R}_+.$$

Let us push-forward the function τ to S by φ_α,

$$\tau_\alpha := \tau \circ \varphi_\alpha^{-1}.$$



Figure 9: Function a_t, $t = 0.3$

2.7.2. (Local wrinkling) *For sufficiently big N and sufficiently small γ the map $I_S^{\tau_\alpha} \circ g : S \to \mathbb{R}^{n+1}$ is a wrinkled embedding. Here $g : S \to S$ is an appropriate smooth homeomorphism, which makes the composition smooth.*

Remark. Of course, $I_S^{\tau_\alpha}(S) = (I_S^{\tau_\alpha} \circ g)(S)$. However, the map $I_S^{\tau_\alpha}$ itself is not smooth. Using the property (b) of the function a_t one can write down the reparameterization g explicitly.

Let $D^1_{\mathbb{Q}}$ be the set of rational points in D^1. We can choose parameterizations φ_α is such a way that the images $\varphi_\alpha(D^{n-1} \times D^1_{\mathbb{Q}}) \subset S$ are pairwise disjoint. Then for a sufficiently large N the images

$$\varphi_\alpha(\bigcup_{-N}^{N}[x_k - \widetilde{\sigma}, x_k + \widetilde{\sigma}]) \subset S,$$

where $x_k = \frac{2k}{2N+1}, \widetilde{\sigma} = \frac{4\sigma}{2N+1}$, are pairwise disjoint as well.

Choose a partition of unity $\sum_1^K \eta_\alpha = 1$ on $\widetilde{S}$ realized by functions η_α supported in $\operatorname{Int} C_\alpha$ and such that $\eta_\alpha|_{\widetilde{C}_\alpha} > 0$. Finally, for a sufficiently large N and sufficiently small $\gamma > 0$, the map $I_S^\Psi \circ g$, where

$$\Psi := \sum_{\alpha=1}^{K} \eta_\alpha \tau_\alpha,$$



Figure 10: Function $\mathcal{W}_{\sigma,1},\ x \in [-\frac{1}{2}, \frac{1}{2}],\ \sigma \simeq 0.02$

is the required approximation of i_S by an almost horizontal wrinkled embedding. Here, like in 2.7.2, $g : S \to S$ is an appropriate reparameterization. □

We will need the following *parametric* version of Lemma 2.7.1:

2.7.3. (Approximation of a family of embedded hypersurfaces by a family of almost horizontal wrinkled embeddings) *Let $S_t \subset \mathbb{R}^{n+1}$, $t \in I$, be a family of oriented quasi-graphical hypersurfaces, such that S_t is almost horizontal for $t = 0$ and S_t is almost horizontal near the boundary ∂S_t for all $t \in I$. Then there exists a C^0-approximation of the family of embeddings $i_{S_t} : S_t \hookrightarrow \mathbb{R}^{n+1}$ by a family of almost horizontal wrinkled embeddings $f_t : S_t \to \mathbb{R}^{n+1}$ of depth $\leqslant 1$, such that f_t coincide with i_{S_t} for $t = 0$ and f_t coincide with i_{S_t} near ∂S for all $t \in I$.*

In order to prove this version, we need to do the following modification in the previous proof. The family S_t, $t \in I$, can be considered as a fibered over I quasi-graphical hypersurface S in $I \times \mathbb{R}^{n+1}$. Then, all above constructions should be done in the fibered category. In particular, instead of the cylinder $D^n \times D^1$ we need to use the fibered cylinder $I \times D^{n-1} \times D^1$ for the parameterization of the fibered cylindrical sets.
The hypersurface S is almost horizontal everywhere near the boundary ∂S except the right side, $S_1 \subset \partial S$. Hence, in order to use the above scheme,

we will work with the doubled, fibered over $[0,2]$ family $S_t, t \in [0,2]]$, where $S_t = S_{2-t}$ for $t \in [1,2]$, while making all the above constructions equivariant with respect to the involution $t \mapsto 2-t$, $t \in [0,2]$.

2.8 Proof of Theorem 2.5.2

In order to get rid of non-essential details, we will consider only the case $(W, g_W) = (\mathbb{R}^q, dx_1^2 + ... + dx_m^2)$. The proof can be easily adjusted to the general case.

Working with a triangulation Δ of the manifold V, we always assume that $S' \subset S = \Sigma(f_0)$ is contained in the $(n-2)$-skeleton of Δ and S in contained in its $(n-1)$-skeleton. Given a triangulation Δ of V and a map $G : V \to \mathrm{Gr}_n\mathbb{R}^m$, we will denote by G^Δ the piecewise constant map $G^\Delta : V \to \mathrm{Gr}_n\mathbb{R}^m$, defined on each n-simplex Δ_i^n by the condition that $G^\Delta(v)$ is parallel to $G(v_i)$, where v_i is the barycenter of the simplex Δ_i^n. The map G^Δ is multivalued over the $(n-1)$-skeleton of Δ.

The case $\mathbf{m = n+1}$. We can choose a triangulation Δ of V with sufficiently small simplices (*fine* triangulation), such that for any i the image $f_0(\Delta_i^n)$ is arbitrarily C^1-close to $G_0(v_i)$ and for any t the map G_t^Δ is arbitrarily C^0-close to G_t. In what follows we will approximate the homotopy G_t^Δ instead of G_t, and therefore the approximation $G_t \approx G_t^\Delta$ must have at least the same order as the desired approximation $G_t \approx \mathrm{G}df_t$.

First, we apply Theorems 2.2.1 and 2.6.1 to the $(n-1)$-skeleton Δ^{n-1} and construct a graphical isotopy $\widetilde{f}_t : V \to \mathbb{R}^{n+1}$, $\widetilde{f}_0 = f_0$, such that $\mathrm{G}d\widetilde{f}_t|_{\mathcal{O}p\,\Delta^{n-1}}$ is arbitrarily C^0-close to $G_t|_{\mathcal{O}p\,\Delta^{n-1}}$. For every i and t the hypersurface $\widetilde{f}_t(\Delta_i^n)$, i.e. the image of $f_0(\Delta_i^n)$ under the graphical isotopy, is almost graphical with respect to $\mathrm{G}_0(v_i)$, and hence it is quasi-graphical with respect to the hyperplane $G_t(v_i)$ because the angle between $\mathrm{G}_0(v_i)$ and $G_t(v_i)$ is less then $\pi/4$. Then, $\widetilde{f}_t(\Delta_i^n)$ is almost horizontal with respect to $G_t(v_i)$ near the boundary $\partial\widetilde{f}_t(\Delta_i^n)$. Therefore, over each n-simplex Δ_i^n we can apply Lemma 2.7.3 with $S_t = \widetilde{f}_t(\Delta_i^n)$ and $G_t(v_i)$ as the horizontal hyperplane, i.e. for every t the hyperplane $G_t(v_i)$ plays the role of the horizontal hyperplane $\mathbb{R}^n \subset \mathbb{R}^{n+1}$. Thus we can deform the isotopy of embeddings $\widetilde{f}_t$ to the desired homotopy f_t of the wrinkled embeddings.

The case $\mathbf{m > n+1}$. Let Δ be a fine triangulation of V. For every i the image $f_0(\Delta_i^n)$ is arbitrarily C^1-close to $G_0(v_i)$ and, therefore, we can work

over each simplex Δ^n_i with the projection of the isotopy $\widetilde{f}_t$ to $L_{v_i} \simeq \mathbb{R}^{n+1}$ (the $(n+1)$-dimensional subspace where the rotation G_t goes on) exactly as in the previous case, keeping the coordinates in $L^{\perp}_{v_i}$ unchanged. $\square$

2.9 Fibered case

Let us formulate and discuss the parametric version of Theorem 2.5.1. For a fibration $q : W \to B$ denote by $\mathrm{Gr}_n W_B$ the Grassmannian of n-planes tangent to the fibers of the fibration q.

2.9.1. (Global integrable approximation of fibered tangential rotation) *Let $G_t : V \to \mathrm{Gr}_n W$ be a fibered tangential rotation of a fibered (over B) embedding $i_V : V \hookrightarrow W$. Then there exists a homotopy of fibered wrinkled embeddings $f_t : V \to W$, $f_0 = i_V$, such that the (fibered) homotopy $\mathrm{G}d_B f_t : V \to \mathrm{Gr}_n W_B$ is arbitrarily C^0-close to G_t. If the rotation G_t is fixed on a closed subset $C \subset V$, then the homotopy f_t can be chosen also fixed on C. In particular, if G_t is fixed over a closed subset $B' \subset B$, then the homotopy f_t can be chosen also fixed over B'.*

This theorem follows from

2.9.2. (Global integrable approximation of simple fibered tangential rotation) *Let $G_t : V \to \mathrm{Gr}_n W$ be a simple fibered tangential rotation of a fibered* **wrinkled** *embedding $f_0 : V \to W$. Then there exists a homotopy of fibered wrinkled embeddings $f_t : V \to W$, such that the (fibered) homotopy $\mathrm{G}d_B f_t : V \to \mathrm{Gr}_n W_B$ is arbitrarily C^0-close to G_t.*

Now we need a version of the main lemma 2.7.1 which is parametric with respect to both, time and space:

2.9.3. (Approximation of family of fibered embedded hypersurfaces by family of almost horizontal fibered wrinkled embeddings) *Let $S_t \subset B^k \times \mathbb{R}^{n+1}$, $t \in I$, be a family of fibered over B^k oriented quasi-graphical hypersurfaces, such that S_t is almost horizontal for $t = 0$ and S_t is almost horizontal near the boundary ∂S_t and over ∂B for all $t \in I$. Then there exists a C^0-approximation of the family of fibered embeddings $i_{S_t} : S_t \hookrightarrow B \times \mathbb{R}^{n+1}$ by a family of almost horizontal fibered wrinkled embeddings $f_t : S_t \to \mathbb{R}^{n+1}$ of depth $\leqslant 1$, such that f_t coincides with i_{S_t} for $t = 0$ and f_t coincides with i_{S_t} near ∂S and over ∂B for all $t \in I$.*

The proof of the time-parameric version 2.7.3 of the main lemma can be rewritten for this fibered case. In the proof of Theorem 2.9.2, in order to apply the fibered version of the main lemma, the fine triangulation Δ ought to be transversal to the fibers on $\operatorname{Int}\Delta_i$ for all its simplices $\Delta_i \subset S \setminus \mathcal{O}p\,\partial S$. Let us point that the existence of such a triangulation is non-obvious and was proven in a more general case (for foliation) by W. Thurston in [Th74].

2.10 Double folds

A (smooth) map $f : V^n \to W^m\,,\ n < m\,$, is called *folded embedding*, if

- f is a topological embedding;
- the singularity $\Sigma = \Sigma_f$ of the map f is an $(n-1)$-dimensional submanifold in V;
- near each connected components S of Σ the map f is equivalent to the map
$$\mathcal{O}p_{\,S\times\mathbb{R}^1} S \times 0 \to S \times \mathbb{R}^{m-n+1}$$
given by the formula $(y, z) \to (y, z^2, z^3, 0, ..., 0)$, where $y \in S$.

Thus, folded embeddings have only *two-fold* corners, ike on Fig.1. The submanifolds S will be called the *folds* of the folded embedding f.
We say that a folded embedding f has *spherical double-folds* if all its folds diffeomorphic to the the $(n-1)$-sphere, and organized in pairs (S_0, S_1) which bounds annuli diffeomorphic to $S^{n-1} \times I$. These annuli are allowed to be *nested*, i.e. the annulus bounded by one double fold may contain the annulus associated with another double fold.
Considering families of folded embeddings with spherical double folds we will also allow, similar to the case of wrinkled embeddings, *embryo* double folds, so that a double fold could die or be born during the deformation. The local model for a map $\mathcal{O}p_{\,S\times\mathbb{R}^1} S \times 0 \to S \times \mathbb{R}^{m-n+1}$ near an embryo double fold is given by the formula

$$(y, z) \longmapsto (y, z^3, z^5, 0, ..., 0).$$

Theorem 2.5.1 has an equivalent reformulation for folded embeddings with spherical double folds.

2.10.1. (Global integrable approximation of tangential rotation by folded embeddings with spherical double folds) *Let $G_t : V \to \mathrm{Gr}_n W$ be a tangential rotation of an embedding $i_V : V \hookrightarrow W$. Then there exists a homotopy of folded embeddings with spherical double folds $f_t : V \to W$, $f_0 = i_V$, such that the homotopy $\mathrm{G}df_t : V \to \mathrm{Gr}_n W$ is arbitrarily C^0-close to G_t. If the rotation G_t is fixed on a closed subset $C \subset V$, then the homotopy f_t can be chosen also fixed on C.*

A fibered version of this theorem also holds.

The following surgery construction allows one to deduce 2.10.1 from 2.5.1. Let $f : V^n \to W^m$ be a wrinkled embedding, $n > 1$. One can modify each wrinkle of f by a connected sum construction for cusps (Whitney surgery) such that the resulting map $\widetilde{f} : V^n \to W^m$ will be a folded embedding with *spherical double folds* $S^{n-1} \times S^0$. For maps $V^2 \to W^m$ the construction (in the pre-image) is shown on Fig.11. Next two propositions contain a formal



Figure 11: Whitney surgery (in the pre-image); $n = 2$

description of the construction, see [El72] or [EM98] for more details.

2.10.2. (Preparation for the cusp surgery) *Let*

$$Z(n,q) : \mathcal{O}p_{\,\mathbb{R}^n} S^{n-1} \to \mathbb{R}^m$$

be the standard wrinkled embedding with the wrinkle $S^{n-1} \subset \mathbb{R}^n$. If $n > 1$ then there exists an embedding $h : D^{n-1} \to \mathcal{O}p_{\,\mathbb{R}^n} S^{n-1}$ such that

- *$h(\partial D^{n-1}) = S^{n-2}$, the cusp (three-fold points) of the wrinkle;*
- *$h(D^{n-1} \setminus \mathrm{Int}\, D^{n-1}_{1-\delta}) = (D^{n-1}_{1+\delta} \setminus \mathrm{Int}\, D^{n-1}) \times 0 \subset \mathbb{R}^{n-1} \times \mathbb{R}^1$;*
- *$h(\mathrm{Int}\, D^{n-1})$ does not intersect the wrinkle S^{n-1}.*

2.10.3. *(***Surgery of cusps)** *Let $h : D^{n-1} \to \mathcal{O}p_{\mathbb{R}^n} S^{n-1}$ be an embedding, as in 2.10.2. There exists a C^0-small perturbation of the map $Z(n,m)$ in an arbitrarily small neighborhood of the embedded disk $h(D^{n-1})$ such that the resulting map $\widetilde{Z}(n,m)$ is a folded embedding with two spherical folds.*

The construction also implies that each double fold $S^{n-1} \times S^0$ of the folded embeddings $\widetilde{f}$ bounds an annulus $S^{n-1} \times D^1$ in V.

3 Applications

3.1 Homotopy principle for directed wrinkled embeddings

Using the wrinkled embeddings one can reformulate the h-principle for A-directed embeddings of *open* manifolds (see 4.5.1 in [EM02]) to the case of A-directed wrinkled embeddings of *closed* manifolds.

3.1.1. (A-directed wrinkled embeddings of closed manifolds) *If $A \subset \mathrm{Gr}_n W$ is an open subset and $f_0 : V \hookrightarrow W$ is an embedding whose tangential lift*

$$G_0 = \mathrm{G}df_0 : V \to \mathrm{Gr}_n W$$

is homotopic to a map

$$G_1 : V \to A \subset \mathrm{Gr}_n W\,,$$

then there exists a homotopy of wrinkled embeddings $f_t : V \to W$ such that $f_1 : V \to W$ is an A-directed wrinkled embedding. Such a homotopy can be chosen arbitrarily C^0-close to f_0.

One can formulate 3.1.1 also in the case when A is an open subset in the Grassmannian of *oriented* n-planes in W. In both cases the h-principle holds also in the relative and parametric versions. Then, one can reformulate this h-principle for the folded embeddings with spherical double folds.

3.2 Embeddings into foliations

A. Wrinkled mappings and generalized wrinkled mappings. In Section 4 below we provide for a convenience of the reader the basic definitions

and results of the wrinkling theory from [EM97] and [EM98]. However, in this section we talk about *generalized* wrinkled maps. Let us explain here the difference between the two notions.

A wrinkled mapping $f : V^n \to W^q$, $n \geqslant q$, by definition, is a map whose singularity set consists of (n, q)-*wrinkles* (= *wrinkles*), where each wrinkle is a $(q-1)$-dimensional sphere $S_i \subset V$ of *fold* points divided by an equator of *cusp* points, see Fig. 12. In addition, each wrinkle $S_i \subset V$ is required to



Figure 12: Wrinkle and its image; $n = q = 2$

bound an embedded q-dimensional disk $D_i \subset V$, such that *the restriction of the map f to $D_i \setminus S_i$ is an equidimensional embedding.*



Figure 13: Nested wrinkles in the pre-image (i.e. in V); $n = 2$

In the case of a *generalized wrinkled mapping* f each (n, q)-wrinkle S_i also bounds a disk D_i in V. However, the restriction $f|_{D_i}$ *is not required to be a smooth embedding on $D_i \setminus S_i$*. Instead, this restriction is allowed itself to be wrinkled, see Fig.13. See Section 4.6 below for more details.

B. Main theorem (statement). Let $(W^m, \mathcal{F})$ be a foliated manifold, $\operatorname{codim} \mathcal{F} = q \leqslant n$. We say that the singularities of an embedding $f : V^n \to (W, \mathcal{F})$ with respect to foliation $\mathcal{F}$ are *generalized wrinkles*, if f is a *generalized wrinkled map with respect to* $\mathcal{F}$, see Section 4.6.

3.2.1. (Embeddings into foliations) *Let $\mathcal{F}_p$, $p \in B$, be a family of foliations on W^m, $\operatorname{codim} \mathcal{F}_p = q \leqslant n$, and $f_p : V^n \to (W^m, \mathcal{F}_p)$ be a family of embeddings such that f_p is transversal to $\mathcal{F}_p$ for $p \in \mathcal{O}p\, \partial B$. Suppose, in addition, that there exists a family of tangential rotations $G_p^t : V \to \operatorname{Gr}_n W$, $p \in B$, $t \in I$, such that*

- *G_p^t is constant for $p \in \mathcal{O}p\, \partial B$;*
- *G_p^1 is transversal to the foliation $\mathcal{F}_p$ for all p.*

Then there exists a family of embeddings f_p^t, $p \in B$, $t \in I$, such that

- *$f_p^0 = f_p$ for all p and $f_p^t = f_p$ for $p \in \mathcal{O}p\, \partial B$;*
- *for any p the singularities of the f_p^1 with respect to $\mathcal{F}_p$ are generalized (n, q)-wrinkles and embryos.*

◀ **Remarks. 1.** The family f_p^t is C^0-close to the constant family $\bar{f}_p^t = f_p^0$.
2. The theorem is true also relative to a closed subset $A \subset V$, i.e. in the situation when the embeddings f_p are already transversal to $\mathcal{F}$ on $\mathcal{O}p_{\,V} A$.
3. In a more formal way, one can say that f_p^1 is a *fibered generalized wrinkled map with respect to $\mathcal{F}_p$*. ▶

◀ **Examples. 1.** *Let $f_p : S^{n-1} \times I \to \mathbb{R}^{n+1}$, $p \in D^k$, be a family of embeddings, such that f_p is the standard inclusion $i_{S^{n-1} \times I} : S^{n-1} \times I \hookrightarrow \mathbb{R}^{n+1}$ for $p \in \mathcal{O}p\, \partial D^k$ and $f_p = i_{S^{n-1} \times I}$ near $\partial(S^{n-1} \times I)$ for all p. Then there exists an isotopy of the family f_p which is fixed near $\partial S^{n-1} \times I$ and for $p \in \partial D^k$ and such that the projections of the resulting cylinders to the axis x_{n+1} have only Morse and birth-death type singularities.*
Indeed, the corresponding homotopical condition for the family of normal vector fields is automotically holds here, as it was observed by A. Douady and F. Laudenbach, see [La76] and [EM00]. Similarly, we have
2. *Let $f_p : D^n \to \mathbb{R}^{n+1}$, $p \in D^k$ be a family of embeddings, such that f_p is the standard inclusion $i_{D^n} : D^n \hookrightarrow \mathbb{R}^{n+1}$ for $p \in \mathcal{O}p\, \partial D^k$, and $f_p = i_{D^n}$ near ∂D^n for all p. Then there exists an isotopy of the family f_p, which is fixed near ∂D^n and for $p \in \partial D^k$, such that all the singularities of the projections of the resulting disks on $\mathbb{R}^n$ are generalized wrinkles and embryos.* ▶

Theorem 3.2.1 is proved below in **D**.

C. $\mathcal{F}$-regularization (lemmas). For the canonical form

$$Z(n, n+1) : \mathcal{O}p_{\,\mathbb{R}^n} S^{n-1} \to \mathbb{R}^{n+1}$$

of the wrinkled embedding (see 2.3 A) the *regularizing foliation* is, by definition, the one-dimensional affine foliation in $\mathbb{R}^{n+1}$, parallel to the axis x_{n+1}.

In order to make the situation more transparent and simplify the notation, we formulate our lemmas only in the case $m = n+1$. All statements remain true also for an arbitrary $m \geq n+1$.

3.2.2. *Let $f : V^n \to (W^{n+1}, \mathcal{F})$ be a wrinkled embedding, transversal to the foliation $\mathcal{F}$, $\dim \mathcal{F} = 1$ ($\operatorname{codim} \mathcal{F} = n$). Then there exists a regularization $\widetilde{f}$ of the wrinkled embedding f such that the singularities of the smooth embedding $\widetilde{f}$ with respect to $\mathcal{F}$ are generalized wrinkles.*

Proof. We can chose the canonical coordinates near each wrinkle $f(S_i)$ such that $\mathcal{F}$ will be the regularizing foliation and then apply the standard regularization (see 2.3 B). □

Let $f : V^n \to (W^{n+1}, \mathcal{F})$ be a wrinkled embedding. Denote by $\widehat{V}_i$ an n-dimensional submanifold in W^m, such that $f(S_i) \subset \widehat{V}_i$ and $\widehat{V}_i$ is tangent to $f(\mathcal{O}p\, S_i)$ along $f(S_i)$ (see 2.3 D). Notice that if f is transversal to $\mathcal{F}$ then $\widehat{V}_i$ is transversal to $\mathcal{F}$ near $f(S_i)$.

3.2.3. *Let $f : V^n \to (W^{n+1}, \mathcal{F})$ be a wrinkled embedding, transversal to the foliation $\mathcal{F}$, $\dim \mathcal{F} \geqslant 2$ ($\operatorname{codim} \mathcal{F} \leqslant n-1$). Suppose, in addition, that the restriction $f|_{S_i} : S_i \to \widehat{V}_i$ is transversal to the foliation $\mathcal{F} \cap \widehat{V}_i$ for all i. Then any regularization $\widetilde{f}$ of the wrinkled embedding f (i.e. for any choice of the canonical coordinates) gives us a smooth embedding transversal to $\mathcal{F}$.* □

3.2.4. *Let $f : V^n \to (W^{n+1}, \mathcal{F})$ be a wrinkled embedding, transversal to the foliation $\mathcal{F}$, $\dim \mathcal{F} \geqslant 2$ ($\operatorname{codim} \mathcal{F} \leqslant n-1$). Suppose, in addition, that the singularities of the restriction $f|_{S_i} : S_i \to \widehat{V}_i$ with respect to $\mathcal{F} \cap \widehat{V}_i$ are generalized $(n-1, q)$-wrinkles for all i. Then there exists a regularization $\widetilde{f}$ of the map f such that the singularities of the smooth embedding $\widetilde{f}$ with respect to $\mathcal{F}$ are generalized (n, q)-wrinkles.*

Proof. We can choose the canonical coordinates near each wrinkle $f(S_i)$ in such a way that the one-dimensional regularizing foliation is inscribed into $\mathcal{F}$, and then apply the standard regularization. Such a regularization adds $\pm t^2$ to the canonical form of any generalized $(n-1, q)$-wrinkle and thus transform it to a (n, q)-wrinkle. See Fig. 14. □

Remark. Lemmas 3.2.2- 3.2.4 remain true in the parametric form.



Figure 14: Regularization near the $(1,1)$-wrinkle

D. Main theorem (proof). We consider only a non-parametric situation, i.e. when B is just a point. The proof in the parametric case is similar with a systematic use of the fibered terminology. Thus, we will drop "p" from the notation. Moreover, we will consider only the case $m = n+1$; the proof can be easily rewritten for any $m > n$.

First of all, let us apply Theorem 2.5.1 to the family $G^t : V^n \to \mathrm{Gr}_n W^{n+1}$ and construct a family of wrinkled embeddings $\widehat{f}^t : V^n \to W^{n+1}$, $t \in [0,1]$, transversal to $\mathcal{F}$.

a) *The base of the induction:* $\dim \mathcal{F} = 1$ (the equidimensional case). We regularize the family $\widehat{f}^t$ in such a way that for $\widehat{f}^1$ our regularization is the $\mathcal{F}$-regularization as in Lemma 3.2.2, and get the required family f^t of embeddings where f^1 is a generalized wrinkled map with respect to $\mathcal{F}$.

b) *Wrinkling of wrinkles*: $\dim \mathcal{F} > 1$. Let $\widehat{V}_i \subset W$ be an n-dimensional submanifold, such that $\Sigma_i = \widehat{f}^1(S_i) \subset \widehat{V}_i$ and $\widehat{V}_i$ is tangent to $\widehat{f}^1(V)$ along Σ_i. Here S_i is a wrinkle of the wrinkled embedding $\widehat{f}^1$. The foliation $\mathcal{F}$ is transversal to $\widehat{V}_i$ near Σ_i. Each wrinkle originates from embryo (i.e. from a point) and hence one can rotate the tangent $(n-1)$-planes to $\Sigma_i \subset \widehat{V}_i$ in $\widehat{V}_i$ to a position transversal to the foliation $\mathcal{F} \cap \widehat{V}$. Such a rotation can be approximated near $\Sigma_i' = \widehat{f}^1(S_i')$ by an isotopy of Σ_i in $\widehat{V}_i$. Then, by our *inductional hypothesis* there exists an isotopy of Σ_i in $\widehat{V}_i$, fixed near Σ_i' and such that the singularities of the final embedding with respect to $\mathcal{F} \cap \widehat{V}_i$ are generalized $(n-1,q)$-wrinkles. The resulting isotopies g_i^t, $t \in [1,2]$ (in $\widehat{V}_i$, for all i) can be extended to an isotopy $\widehat{f}^t$, $t \in [1,2]$, of the wrinkled embedding $\widehat{f}^1$. Finally, we regularize the family $\widehat{f}^t$, $t \in [0,2]$, in such a way that for $\widehat{f}^2$ our regularization is the $\mathcal{F}$-regularization as in Lemma 3.2.4, and get the required family f^τ, $\tau \in [0,1], \tau = 2t$, of embeddings such that f^1 is a

generalized wrinkled map with respect to $\mathcal{F}$. □

◄ **Remark.** Using in the above proof Theorem 2.10.1 instead of 2.5.1 we get a version of Theorem 3.2.1 about embeddings with the double fold type tangency singularities to a foliation. ►

3.3 Embeddings into distributions

In this section we sketch a generalization of Theorem 3.2.1 to the case of distributions.

Let ξ be a distribution on a manifold W^m, $\operatorname{codim}\xi = q$. An embedding $f : V^n \to (W^m, \xi)$, $n \geqslant \operatorname{codim}\xi$, is called *transversal to* ξ, if the *reduced differential*

$$d^\xi f : TV \xrightarrow{df} TW \xrightarrow{\pi^\xi} TW/\xi$$

is surjective. The *non-transversality* defines a variety Σ_ξ of the 1-jet space $J^1(V, W)$. For a general non-integrable ξ one cannot define fold and cusp type tangency through normal forms. However, the original Whitney-Thom definition is applicable in this situation as well.
We say that f has at a point $p \in V$ a tangency to ξ of *fold* type if

- $\operatorname{Corank} d^\xi_p = 1$.
- $J^1(f) : V \to J^1(V, W)$ is transverse to Σ;
- $d^\xi_p f|_{T_p\Sigma(f)} : T_p\Sigma(f) \to TW_{f(p)}/\xi$ is injective.

For a codimension 1 distribution ξ defined by a Pfaffian equation $\alpha = 0$, the fold tangency point of an embedding $f : V \to W$ to ξ are isolated non-degenerate zeroes of the induced 1-form $f^*\alpha$ on V.
Similarly, we define a tangency of *cusp* type by the same first two conditions and, in addition by requiring that $d^\xi_p f|_{T_p\Sigma(f)} : T_p\Sigma(f) \to TW_{f(p)}/\xi$ is not injective and the 2-jet section $J^2(f) : V \to J^2(V, W)$ to be transversal to the singularity in $J^2(V, W)$ defined by the non-injectivity condition.
Combining the following observations, one can associate with a co-oriented fold an *index*, as in the case of a foliation.

3.3.1. (Index for fold-type tangency to distribution)

A. *Let $J^2(\xi) \to W^m$ be the bundle of 2-jets of codimension q submanifolds of W tangent to ξ. There exists a section $W \to J^2(\xi)$ which can be uniquely characterized by the following property: for every point $p \in W$ there exists an embedding $\varphi_p : \mathbb{R}^{m-q} \to W$ such that $d_p\varphi_p(\mathbb{R}^{m-q}) = \xi_p$, and for all $t \in \mathbb{R}$, $x \in \mathbb{R}^{m-q}$ we have $\frac{d\varphi_p(tx)}{dt} \in \xi$.*

B. *Let (W, ξ) be as above, and Σ a submanifold of dimension $< \operatorname{codim} \xi$ such that $\pi^\xi : T\Sigma \to TW/\xi$ is injective. Then, given any section $\sigma : \Sigma \to J^2(\xi)$ there exists codimension q foliation $\mathcal{F}$ on $\mathcal{O}p\,\Sigma$ whose 2-jet along Σ is equal to σ.*

C. *Suppose that an embedding $f : V \to W$ has a fold type tangency of index k to a foliation $\mathcal{F}$ along a submanifold $\Sigma \subset W$. Let $\widetilde{\mathcal{F}}$ be another foliation of the same codimension and which have the same 2-jet as $\mathcal{F}$ along Σ. Then $f : V \to W$ has at Σ a fold type tangency to a foliation $\widetilde{\mathcal{F}}$ of the same index k.*

This allows us define generalized (n, q)-wrinkles and embryo type singularities of tangency of an embedding $f : V^n \to (W, \xi)$ to a distribution ξ.
With these definitions Theorem 3.2.1 can be generalized without any changes to the case of an arbitrary distribution ξ instead of a foliation. The inductional proof for the foliation case presented in Section 3.2 works in this more general with a few additional remarks. Namely, one note that the base of the induction, i.e. the case when $\dim \xi = 1$ is the same in this case, as 1-dimensional distributions are integrable. When applying in the inductional step the standard regularization from 2.3 B, one need to choose the regularizing function $\widetilde{u}(y, z)$ sufficiently C^1-close to $u(y, z)$. With this modification the proof goes through as is.

4 Appendix: Wrinkling

We recall here, for a convenience of the reader, some definitions from [EM97] and [EM98] and introduce the notions of *generalized wrinkles* and *generalized wrinkled maps*. We also formulate here some *results* from [EM97] and [EM98] though we do not use these theorems in the paper.

4.1 Folds and cusps

Let V and W be smooth manifolds of dimensions n and q, respectively, and $n \geq q$. For a smooth map $f : V \to W$ we will denote by $\Sigma(f)$ the set of its singular points, i.e.

$$\Sigma(f) = \{p \in V, \ \text{ rank } d_p f < q\} \ .$$

A point $p \in \Sigma(f)$ is called a *fold* type singularity or a *fold* of index s if near the point p the map f is equivalent to the map

$$\mathbb{R}^{q-1} \times \mathbb{R}^{n-q+1} \to \mathbb{R}^{q-1} \times \mathbb{R}^1$$

given by the formula

$$(y, x) \mapsto \left(y, \ -\sum_1^s x_i^2 + \sum_{s+1}^{n-q+1} x_j^2 \right) \tag{5}$$

where $x = (x_1, \ldots, x_{n-q+1}) \in \mathbb{R}^{n-q+1}$ and $y = (y_1, ..., y_{q-1}) \in \mathbb{R}^{q-1}$. For $W = \mathbb{R}^1$ this is just a nondegenerate index s critical point of the function $f : V \to \mathbb{R}^1$.

Let $q > 1$. A point $p \in \Sigma(f)$ is called a *cusp* type singularity or a *cusp* of index $s + \frac{1}{2}$ if near the point p the map f is equivalent to the map

$$\mathbb{R}^{q-1} \times \mathbb{R}^1 \times \mathbb{R}^{n-q} \to \mathbb{R}^{q-1} \times \mathbb{R}^1$$

given by the formula

$$(y, z, x) \mapsto \left(y, z^3 + 3y_1 z - \sum_1^s x_i^2 + \sum_{s+1}^{n-q} x_j^2 \right) \tag{6}$$

where $x = (x_1, \ldots, x_{n-q}) \in \mathbb{R}^{n-q}$, $z \in \mathbb{R}^1$, $y = (y_1, \ldots, y_{q-1}) \in \mathbb{R}^{q-1}$.

For $q \geq 1$ a point $p \in \Sigma(f)$ is called an *embryo type* singularity or an *embryo* of index $s + \frac{1}{2}$ if f is equivalent near p to the map

$$\mathbb{R}^{q-1} \times \mathbb{R}^1 \times \mathbb{R}^{n-q} \to \mathbb{R}^{q-1} \times \mathbb{R}^1$$

given by the formula

$$(y, z, x) \mapsto \left(y, z^3 + 3|y|^2 z - \sum_1^s x_i^2 + \sum_{s+1}^{n-q} x_j^2 \right) \tag{7}$$

where $x \in \mathbb{R}^{n-q},\ y \in \mathbb{R}^{q-1},\ z \in \mathbb{R}^1,\ |y|^2 = \sum_1^{q-1} y_i^2$.

Notice that folds and cusps are stable singularities for individual maps, while embryos are stable singularities only for 1-parametric families of mappings. For a generic perturbation of an individual map embryos either disappear or give birth to wrinkles which we consider in the next section.

4.2 Wrinkles and wrinkled mappings

Consider the map

$$w(n,q,s) : \mathbb{R}^{q-1} \times \mathbb{R}^1 \times \mathbb{R}^{n-q} \to \mathbb{R}^{q-1} \times \mathbb{R}^1$$

given by the formula

$$(y,z,x) \mapsto \left(y, z^3 + 3(|y|^2 - 1)z - \sum_1^s x_i^2 + \sum_{s+1}^{n-q} x_j^2 \right), \tag{8}$$

where $y \in \mathbb{R}^{q-1},\ z \in \mathbb{R}^1,\ x \in \mathbb{R}^{n-q}$ and $|y|^2 = \sum_1^{q-1} y_i^2$.

Notice that the singularity $\Sigma(w(n,q,s))$ is the $(q-1)$-dimensional sphere

$$S^{q-1} = S^{q-1} \times 0 \subset \mathbb{R}^q \times \mathbb{R}^{n-q}.$$

Its equator $\{|y| = 1, z = 0, x = 0\} \subset \Sigma(w(n,q,s))$ consists of cusp points of index $s + \frac{1}{2}$. The upper hemisphere $\Sigma(w) \cap \{z > 0\}$ consists of folds of index s and the lower one $\Sigma(w) \cap \{z < 0\}$ consists of folds of index $s+1$. Also it is useful to notice that the restrictions of the map $w(n,q,s)$ to subspaces $y_1 = t$, viewed as maps $\mathbb{R}^{n-1} \to \mathbb{R}^{q-1}$, are non-singular maps for $|t| > 1$, equivalent to $w(n-1, q-1, s)$ for $|t| < 1$ and to embryos for $t = \pm 1$.

Although the differential $dw(n,q,s) : T(\mathbb{R}^n) \to T(\mathbb{R}^q)$ degenerates at points of $\Sigma(w)$, it can be canonically *regularized* over $\mathcal{O}p_{\mathbb{R}^n} D^q$, an open neighborhood of the disk $D^q = D^q \times 0 \subset \mathbb{R}^q \times \mathbb{R}^{n-q}$. Namely, we can change the element $3(z^2+|y|^2-1)$ in the Jacobi matrix of $w(n,q,s)$ by a function γ which coincides with $3(z^2 + |y|^2 - 1)$ on $\mathbb{R}^n \setminus \mathcal{O}p_{\mathbb{R}^n} D^q$ and does not vanish along the q-dimensional subspace $\{x = 0\} = \mathbb{R}^q \times \mathbf{0} \subset \mathbb{R}^n$. The new bundle map $\mathcal{R}(dw) : T(\mathbb{R}^n) \to T(\mathbb{R}^q)$ provides a homotopically canonical extension of the

map $dw : T(\mathbb{R}^n \setminus \mathcal{O}p_{\mathbb{R}^n} D^q) \to T(\mathbb{R}^q)$ to an epimorphism (fiberwise surjective bundle map) $T(\mathbb{R}^n) \to T(\mathbb{R}^q)$. We call $\mathcal{R}(dw)$ the *regularized differential* of the map $w(n, q, s)$.

A smooth map $f : V^n \to W^q$, $n \geqslant q$, is called *wrinkled*, if any connected component S_i of the singularity $\Sigma(f)$ is diffeomorphic to the standard $(q-1)$-dimensional sphere S^{q-1} and bounds in V a q-dimensional disk D_i, such that the map $f|_{\mathcal{O}p\, D_i}$ is equivalent to the map $w(n, q, s)|_{\mathcal{O}p_{\mathbb{R}^n} D^q}$. The spheres S_i and its images $f(S_i)$ are called *wrinkles* of the wrinkled mapping f. The differential $df : T(V) \to T(W)$ of the wrinkled map f can be regularized (near each wrinkle and hence globally) to obtain an epimorphism $\mathcal{R}(df) : T(V) \to T(W)$.

4.3 Fibered wrinkles and fibered wrinkled mappings

For any integer $k > 0$ the map $w(k + n, q, s)$ can be considered as a fibered map over $\mathbb{R}^k \times \mathbf{0} \subset \mathbb{R}^{k+n}$. We shall refer to this fibered map as $w_k(k + n, q, s)$. The regularized differential $\mathcal{R}(dw_k(k+n, q, s))$ is a fibered (over $\mathbb{R}^k$) epimorphism

$$\mathbb{R}^k \times T(\mathbb{R}^{q-1} \times \mathbb{R}^1 \times \mathbb{R}^{n-q}) \xrightarrow{\mathcal{R}(dw_k(k+n,q,s))} \mathbb{R}^k \times T(\mathbb{R}^{q-1} \times \mathbb{R}^1)$$

A fibered (over B) map $f : V^{k+n} \to W^{k+q}$, $n \geqslant q$, is called *fibered wrinkled*, if any connected component S_i of the singularity $\Sigma(f)$ is diffeomorphic to the standard $(k + q - 1)$-dimensional sphere S^{k+q-1} and bounds in V a $(k + q)$-dimensional disk D_i, such that the fibered map $f|_{\mathcal{O}p\, D_i}$ is equivalent to the fibered map $w_k(k+n, q, s)|_{\mathcal{O}p_{\mathbb{R}^{k+n}} D^q}$. The spheres S_i and its images $f(S_i)$ are called *fibered wrinkles* of the fibered wrinkled mapping f. The restrictions of a fibered wrinkled map to a fiber may have, in addition to wrinkles, emdryos singularities. For a fibered wrinkled map $f : V \to W$ one can define its regularized differential which is a fibered (over B) epimorphism $\mathcal{R}(d_B f) : T_B V \to T_B W$.

4.4 Main theorems about wrinkled mappings

The following Theorem 4.4.1 and its parametric version 4.4.2 are the main results of our paper [EM97]:

4.4.1 (Wrinkled mappings)**.** *Let $F : T(V) \to T(W)$ be an epimorphism which covers a map $f : V \to W$. Suppose that f is a submersion on a neighborhood of a closed subset $K \subset V$, and F coincides with df over that neighborhood. Then there exists a wrinkled map $g : V \to W$ which coincides with f near K and such that $\mathcal{R}(dg)$ and F are homotopic rel. $T(V)|_K$. Moreover, the map g can be chosen arbitrarily C^0-close to f, and his wrinkles can be made arbitrarily small.*

4.4.2 (Fibered wrinkled mappings)**.** *Let $f : V \to W$ be a fibered over B map covered by a fibered epimorphism $F : T_B(V) \to T_B(W)$. Suppose that F coincides with df near a closed subset $K \subset V$ (in particular, f is a fibered submersion near K), then there exists a fibered wrinkled map $g : V \to W$ which extends f from a neighborhood of K, and such that the fibered epimorphisms $\mathcal{R}(dg)$ and F are homotopic rel. $T_B(M)|_K$. Moreover, the map g can be chosen arbitrarily C^0-close to f, and his wrinkles can be made arbitrarily small.*

4.5 Wrinkled mappings into foliations

Here we formulate a slightly strengthened version of Theorems 4.4.1 and 4.4.2. Let us start with some definitions.

Let $\mathcal{F}$ be a foliation on a manifold W, $\operatorname{codim} \mathcal{F} = q$. A map $f : V \to W$ is called *transversal to* $\mathcal{F}$, if the reduced differential

$$TV \xrightarrow{df} TW \xrightarrow{\pi_{\mathcal{F}}} \nu(\mathcal{F})$$

is an epimorphism. Here $\nu(\mathcal{F}) = TW/\tau(\mathcal{F})$ is the normal bundle of the foliation $\mathcal{F}$.

An open subset $U \subset W$ is called *elementary* (with respect to $\mathcal{F}$), if $\mathcal{F}|_U$ is generated by a submersion $p_{_U} : U \to \mathbb{R}^q$. An open subset $U \subset V$ is called *small* (with respect to f and $\mathcal{F}$), if $f(U)$ is contained in an elementary subset U' of W. A map $f : V \to W$ is called $\mathcal{F}^\perp$*-wrinkled*, or *wrinkled with respect to* $\mathcal{F}$, if there exist disjoint small subsets $U_1, ...U_l \subset V$ such that $f|_{V \setminus (U_1 \cup ... \cup U_l)}$ is transversal to $\mathcal{F}$ and for each $i = 1, ..., l$ the composition

$$U_i \xrightarrow{f|_{U_i}} U_i' \xrightarrow{p_{U_i'}} \mathbb{R}^q$$

(where $U_i' \supset f(U_i)$ is an elementary subset of W), is a wrinkled map. In order to get the *regularized* reduced differential

$$\mathcal{R}(\pi_{\mathcal{F}} \circ df) : TM \to \nu(\mathcal{F})$$

of the $\mathcal{F}^{\perp}$-wrinkled map f, we regularize the differential of each wrinkled map $w_i = p_{_{U_i'}} \circ f|_{U_i}$ as in Sections 4.2 and then set

$$\mathcal{R}(\pi_{\mathcal{F}} \circ df|_{U_i}) = [dp_{U_i'}|_{\nu(\mathcal{F})}]^{-1} \circ \mathcal{R}(dw_i).$$

Similarly to Section 4.3 we can define a *fibered* $\mathcal{F}^{\perp}$-wrinkled map $f : V \to Q$, where the foliation $\mathcal{F}$ on W is fibered over the same base B. Finally we define, in a usual way, the *regularization*

$$\mathcal{R}(\pi_{\mathcal{F}} \circ d_B f) : T_B V \to \nu_B(\mathcal{F})$$

of the fibered reduced differential

$$T_B V \xrightarrow{d_B f} TW \xrightarrow{\pi_{\mathcal{F}}} \nu_B(\mathcal{F}).$$

4.5.1. (Wrinkled mappings of manifolds into foliations) *Let $\mathcal{F}$ be a foliation on a manifold W and let $F : TV \to \nu(\mathcal{F})$ be an epimorphism which covers a map $f : V \to W$. Suppose that f is transversal to $\mathcal{F}$ in a neighborhood of closed subset K of V, and F coincides with the reduced differential $\pi_{\mathcal{F}} \circ df$ over that neighborhood. Then there exists a $\mathcal{F}^{\perp}$-wrinkled map $g : V \to W$ such that g coincides with f near K, and $\mathcal{R}(\pi_{\mathcal{F}} \circ dg)$ is homotopic to F relative to $TV|_K$.*

Proof. Take a triangulation of the manifold V by *small* simplices. First we use Gromov-Phillips' theorem (see [Ph67], [Gr86] or [EM02]) to approximate f near the $(n-1)$-skeleton of the triangulation by a map transversal to $\mathcal{F}$. Then, using Theorem 4.4.1, for a neighborhood U_i of every n-simplex σ_i and an elementary set $U_i' \supset f(U_i)$ we can approximate the map $p_{U_i'} \circ f|_{U_i}$ by a wrinkled map. This approximation can be realized by a deformation of the map f, keeping it fixed on a closed subset of U_i, where f was already previously defined. This process produces the desired $\mathcal{F}^{\perp}$-wrinkled map. $\square$

Similarly, Theorem 4.4.2 can be generalized to the following fibered version of Theorem 4.5.1.

4.5.2. (Fibered wrinkled mappings of manifolds into foliations) *Let $f : V \to W$ be a fibered over B map, $\mathcal{F}$ be a fibered over B foliation on W and let $F : T_B(V) \to \nu_B(\mathcal{F})$ be a fibered epimorphism which covers f. Suppose that f is fiberwise transversal to $\mathcal{F}$ near a closed subset $K \subset V$, and F coincides with fibered reduced differential*

$$T_B V \xrightarrow{d_B f} TW \xrightarrow{\pi_{\mathcal{F}}} \nu_B(\mathcal{F}).$$

near K. Then there exists a fibered $\mathcal{F}^{\perp}$-wrinkled map $g : V \to W$ which extends f from a neighborhood of K, and such that the fibered epimorphisms $\mathcal{R}(\pi_{\mathcal{F}} \circ d_B g)$ and F are homotopic rel. $T_B(V)|_K$.

4.6 Generalized wrinkled mappings

A smooth map $f : V^n \to W^q$, $n \geqslant q$, is called *generalized wrinkled*, if any connected components S_i of the singularity $\Sigma = \Sigma_f$ is diffeomorphic to the standard $(q-1)$-dimensional sphere S^{q-1}, which bounds in V a q-dimensional disk D_i, and for each such sphere the map $f|_{\mathcal{O}p\, S_i}$ is equivalent to the map $w(n,q,s)|_{\mathcal{O}p_{\mathbb{R}^n} S^{q-1}}$. The spheres S_i and its images $f(S_i)$ are called *generalized wrinkles* of the generalized wrinkled mapping f.

A fibered map $f : V^{k+n} \to W^{k+q}$, $n \geqslant q$, is called *generalized fibered wrinkled map*, if any connected components S_i of the singularity $\Sigma = \Sigma_f$ is diffeomorphic to the standard $(k+q-1)$-dimensional sphere S^{k+q-1}, which bounds in V a $(k+q)$-dimensional disk D_i, and for each such sphere the fibered map $f|_{\mathcal{O}p\, S_i}$ is equivalent to the fibered map $w_k(k+n,q,s)|_{\mathcal{O}p_{\mathbb{R}^{k+n}} S^{q-1}}$. The spheres S_i and its images $f(S_i)$ are called *generalized fibered wrinkles* of the generalized wrinkled mapping f.

Let $\mathcal{F}$ be a foliation on a manifold W, $\operatorname{codim} \mathcal{F} = q$. A map $f : V \to W$ is called *generalized wrinkled with respect to $\mathcal{F}$*, if there exist disjoint small subsets $U_1, ...U_l \subset V$ such that $f|_{V \setminus (U_1 \cup ... \cup U_l)}$ is transversal to $\mathcal{F}$ and for each $i = 1, ..., l$ the composition

$$U_i \xrightarrow{f|_{U_i}} U_i' \xrightarrow{p_{U_i'}} \mathbb{R}^q$$

(where $U_i' \supset f(U_i)$ is an elementary subset of W), is a generalized wrinkled map.

References

[Ce68] J. Cerf, *Sur les diffeomorphismes de la sphere de dimension trois* ($\Gamma_4 = 0$), Lecture Notes in Mathematics, No. 53. Springer-Verlag, Berlin-New York 1968.

[El72] Y. Eliashberg, *Surgery of singularities of smooth maps*, Izv. Akad. Nauk SSSR Ser. Mat., **36**(1972), 1321–1347.

[EM97] Y. Eliashberg and N. Mishachev, *Wrinkling of smooth mappings and its applications - I*, Invent. Math., **130**(1997), 345–369.

[EM00] Y. Eliashberg and N. Mishachev, *Wrinkling of smooth mappings - II. Wrinkling of embeddings and K.Igusa's theorem,* Topology, **39**(2000), 711-732.

[EM98] Y. Eliashberg and N. Mishachev, *Wrinkling of smooth mappings - III. Foliation of codimension greater than one,* Topol. Methods in Nonlinear Analysis, **11**(1998), 321-350.

[EM02] Y. Eliashberg and N. Mishachev, *Introduction to the h-principle,* AMS, Graduate Studies in Mathematics, v.48, 2002.

[Gr73] M. Gromov, *Convex integration of partial differential relations*, Izv. Akad. Nauk SSSR Ser. Mat., **37**(1973), 329–343.

[Gr86] M. Gromov, *Partial differential relations*, Springer-Verlag, 1986.

[Ha83] A. Hatcher, *A proof of a Smale conjecture,* $\mathrm{Diff}(S^3) \simeq \mathrm{O}(4)$, Ann. of Math., (2) 117 (1983), no. 3, 553–607.

[Ig84] K. Igusa, *Higher singularities are unnecessary*, Annals of Math., **119**(1984), 1–58.

[La76] F. Laudenbach, *Formes différentielles de degré 1 non singulières: classes d'homotopie de leurs noyaux,* Comment. Math. Helvet., **51**(1976), 447–464.

[Ph67] A. Phillips, *Submersions of open manifolds*, Topology **6**(1967), 171–206.

[RS01] C. Rourke and B. Sanderson, *The compression theorem - I,II*, Geom. and Topology, **5**(2001), 399–429, 431–440.

[RS03] C. Rourke and B. Sanderson, *The compression theorem III: applications*, Alg. and Geom. Topology, 3, (2003), 857–872.

[Sp02] D. Spring, *Directed embeddings and the simplification of singularities*, Comm. in Cont. Math., **4**(2002), 107–144.

[Sp05] D. Spring, *Directed embeddings of closed manifolds*, Comm. in Cont. Math., **7**(2005), 707–725.

[Th74] W. Thurston, *The theory of foliations of codimension greater than one*, Comm. Math. Helvet., **49**(1974), 214–231